

# Alpha Technology: Scientific References

## General

Bossé R, Illy C, Elands J, and Chelsky D

Miniaturizing screening: How low can we go today?  
Drug Discov Today HTS supplement 2000; 42-47

Eglen RM, Reisine T, Roby P, et al

The use of AlphaScreen technology in HTS: current status.  
Curr Chem Genom. 2008; 1:2-10

Sterling J, Bojanic D, Eglen RM, Heyse S, and Strulovici B

Current trends in high-throughput screening.  
Assay Drug Dev Technol. 2008; 6(4):491-504

Ullman EF, Kirakossian H, Switchenko AC, Ishkanian J, Ericson M, Wartchow C, Pirio M, Pease J, Irvin B, Singh S, Singh R, Patel R, Daffon A, Davalian D, Skold C, Kuran N, and Wagner D

Luminescent oxygen channeling assay (LOCI): sensitive, broadly applicable homogeneous immunoassay method.  
Clin Chem. 1996; 42(9):1518-1526

Wu X, Sils MA, and Zhang JH

Further comparison of primary hit identification by different assay technologies and effects of assay measurement variability.  
J Biomol Screen. 2005 10(6):581-589

## Antibody Applications

Lazar GA, Dang W, Karki S, Vafa O, Peng JS, Hyun L, Chan C, Chung HS, Eivazi A, Yoder SC, Vielmetter J, Carmichael DF, Hayes RJ, and Dahiyat BI

Engineered antibody Fc variants with enhanced effector function.

PNAS 2006; 103(11):4005-4010

## Biomarkers

Beasley JR, Swanson R, Yang J, and Dunn D

Conversion of ELISA to no-wash technologies.  
Poster: SBS 2007 Conference, Montreal

Bielefeld-Sevigny M

AlphaLISA immunoassay platform – the “no-wash” high-throughput alternative to ELISA.

Assay Drug Dev Technol. 2009; 7(1):90-92

Cauchon E, Liu S, Percival MD, Rowland SE, Xu D, Binkert C, Strickner P, and Falgoutyret JP

Development of a homogeneous immunoassay for the detection of angiotensin I in plasma using AlphaLISA acceptor beads technology.

Anal Biochem. 2009; 388(1):134-139

Lai MT, Munshi V, Touch S, Tynebor RM, Tucker TJ, McKenna PM, Williams TM, DiStefano DJ, Hazuda DJ, and Miller MD

Antiviral activity of MK-4965, a novel nonnucleoside reverse transcriptase inhibitor.

Antimicrob Agents Ch. 2009; 53(6):2424-2431

Majercak J, Ray WJ, Espeseth A, Simon A, Shi XP, Wolffe C, Getty K, Marine S, Stec E, Ferrer M, Strulovici B, Bartz S, Gates A, Xu M, Huang Q, Ma L, Shughrue P, Burchard J, Colussi D, Pietrak B, Kahana J, Behr D, Rosahl T, Shearman M, Hazuda D, Sachs AB, Koblan KS, Seabrook GR, and Stone DJ

LRRMT3 promotes processing of amyloid-precursor protein by BACE1 and is a positional candidate gene for late-onset Alzheimer's disease.  
PNAS 2006; 103(47):17967-17972

McGiven JA, Sawyer J, Perrett LL, Brew SD, Commander NJ, Fisher A, McLarnon S, Harper K, and Stack JA

A new homogeneous assay for high throughput serological diagnosis of brucellosis in ruminants.

J Immunol Methods. 2008; 337(1):7-15

Poulsen F and Jensen KB

A luminescent oxygen channeling immunoassay for the determination of insulin in human plasma.

J Biomol Screen. 2007; 12(2):240-247

Sehr P, Pawlita M, and Lewis J

Evaluation of different glutathione S-transferase-tagged protein captures for screening E6/E6AP interaction inhibitors using AlphaScreen.

J Biomol Screen. 2007; 12(4):560-567

Szekeres PG, Leong K, Day TA, Kingston AE, and Karran EH

Development of homogeneous 384-well high-throughput screening assays for Abeta1-40 and Abeta1-42 using AlphaScreen technology.

J Biomol Screen. 2008; 13(2):101-111

Wilson J, Rossi CP, Carboni S, Fremaux C, Perrin D, Soto C, Kosco Vilbois M, and Scheer A

A homogeneous 384-well high-throughput binding assay for a TNF receptor using AlphaScreen technology.

J Biomol Screen. 2003; 8(5):522-532

## DNA Quantification

Patel R, Pollner R, de Keczer S, Pease J, Pirio M, DeChene N, Daffon A, and Rose S

Quantification of DNA using the luminescent oxygen channeling assay.  
Clin Chem. 2000; 46(9):1471-1477

## DNA:Protein Interactions

Hornung V, Ablasser A, Charrel-Dennis M, Bauernfeind F, Horvath G, Caffrey DR, Latz E, and Fitzgerald KA

AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC.

Nature. 2009; 458(7237):514-518

## GPCRs

Avlani VA, Gregory KJ, Morton CJ, Parker MW, Sexton PM, and Christopoulos A

Critical role for the second extracellular loop in the binding of both orthosteric and allosteric G protein-coupled receptor ligands.

J Biol Chem. 2007; 282(35):25677-25686

Bösel I, Römler H, Hermsdorf T, Thor D, Busch W, Schulz A, and Schöneberg T

Involvement of the V2 vasopressin receptor in adaptation to limited water supply.

PLoS One. 2009; 4(5):e5573

Caruso ME, Jenna S, Beaulne S, Lee EH, Bergeron A, Chauve C, Roby P, Rual JF, Hill DE, Vidal M, Bossé R, and Chevet E

Biochemical clustering of monomeric GTPases of the Ras superfamily.

Mol Cell Proteomics. 2005; 4(7):936-944

Crouch MF and Osmond RIW

New strategies in drug discovery for GPCRs: high throughput detection of cellular ERK phosphorylation.

Comb Chem High Throughput Screen. 2008; 11, 344-356

- Elster L, Elling C, and Heding A  
Bioluminescence resonance energy transfer as a screening assay: focus on partial and inverse agonism.  
J Biomol Screen. 2007; 12(1):41-49
- Ferguson GM, Valant C, Horne J, Figler H, Flynn BL, Linden J, Chalmers DK, Sexton PM, Christopoulos A, and Scammells PJ  
2-aminothienopyridazines as novel adenosine A1 receptor allosteric modulators and antagonists.  
J Med Chem. 2008; 51(19):6165-6172
- Gabriel D, Vernier M, Pfeifer MJ, Dasen B, Tenaillon L, and Bouhelal R  
High-throughput screening technologies for direct cyclic AMP measurement.  
Assay Drug Dev Technol. 2003; 1(2):291-303
- Gray A, Olsson H, Batty IH, Priganica L, and Peter Downes C  
Nonradioactive methods for the assay of phosphoinositide 3-kinases and phosphoinositide phosphatases and selective detection of signaling lipids in cell and tissue extracts.  
Anal Biochem. 2003; 313(2):234-245
- Hay DL  
Measurement of phosphorylated extracellular signal-regulated kinase 1 and 2 in an undergraduate teaching laboratory with AlphaScreen technology.  
Sci Signal. 2009; 2(62):tr1
- Heinrich JN, Brennan J, Lai MH, Sullivan K, Hornby G, Popiolek M, Jiang L-X, Pausch MH, Stack G, Marquis KL, and Andree TH  
Aplindore (DAB-452), a high-affinity selective dopamine D2 receptor partial agonist.  
Eur J Pharmacol. 2006; 552(1-3):36-45
- Lee HJ, Mun H-C, Lewis NC, Crouch MF, Culverston EL, Mason RS, and Conigrave AD  
Allosteric activation of the extracellular Ca<sup>2+</sup>-sensing receptor by L-amino acids enhances ERK1/2 phosphorylation.  
Biochem J. 2007; 404(1):141-149
- Leroy D, Missotten M, Waltzinger C, Martin T, and Scheer A  
G protein-coupled receptor-mediated ERK1/2 phosphorylation: towards a generic sensor for GPCR activation.  
J Recept Signal Transduct Res. 2007; 27(1):83-97
- May LT, Avlani VA, Langmead CJ, Herdon HJ, Wood MD, Sexton PM, and Christopoulos A  
Structure-function studies of allosteric agonism at M2 muscarinic acetylcholine receptors.  
Mol Pharmacol. 2007; 72(2):463-476
- Morfis M, Tilakaratne N, Furness SGB, Christopoulos C, Werry TD, Christopoulos A, and Sexton PM  
Receptor activity modifying proteins differentially modulate the G protein-coupling efficiency of amylin receptors.  
Endocrinology. 2008; 149(11):5423-5431
- Nawaratne V, Leach K, Suratman N, Loiacono RE, Felder CC, Armbruster BN, Roth BL, Sexton PM, and Christopoulos A  
New insights into the function of M4 muscarinic acetylcholine receptors gained using a novel allosteric modulator and a DREADD (Designer Receptor Activated by a Designer Drug).  
Mol Pharmacol. 2008; 74(4):1119-1131
- Osmond RIW, Sheehan A, Borowicz R, Barnett E, Harvey G, Turner C, Brown A, Crouch MF, and Dyer AR  
GPCR screening via ERK1/2: a novel platform for screening G protein-coupled receptors.  
J Biomol Screen. 2007; 10, 730-737
- Pioszak AA and Xu HE  
Molecular recognition of parathyroid hormone by its G protein-coupled receptor.  
Proc Natl Acad Sci U S A. 2008; 105(13):5034-5039
- Pioszak AA, Parker NR, Suino-Powell K, and Xu HE  
Molecular recognition of corticotropin-releasing factor by its G-protein-coupled receptor CRFR1.  
J Biol Chem. 2008; 283(47):32900-32912
- Siehler S  
Cell-based assays in GPCR drug discovery.  
Biotechnol J. 2008; 3(4):471-483. Review
- Tang Y, Zhou L, Gunnet JW, Wines PG, Cryan EV, and Demarest KT  
Enhancement of arachidonic acid signaling pathway by nicotinic acid receptor HM74A.  
Biochem Biophys Res Commun. 2006; 345(1):29-37
- Valant C, Gregory KJ, Hall NE, Scammells PJ, Lew MJ, Sexton PM, and Christopoulos A  
A novel mechanism of G protein-coupled receptor functional selectivity: muscarinic partial agonist McN-A-343 as a bitopic orthosteric/allosteric ligand.  
J Biol Chem. 2008; 283(43):29312-29321
- Van der Westhuizen ET, Werry TD, Sexton PM, and Summers RJ  
The relaxin family peptide receptor 3 activates extracellular signal-regulated kinase 1/2 through a protein kinase C-dependent mechanism.  
Mol Pharmacol. 2007; 71(6):1618-1629
- Werry TD, Stewart GD, Crouch MF, Watts A, Sexton PM, and Christopoulos A  
Pharmacology of 5HT<sub>2C</sub> receptor-mediated ERK1/2 phosphorylation: agonist-specific activation pathways and the impact of RNA editing.  
Biochem Pharmacol. 2008; 76(10):1276-1287
- Williams C  
cAMP detection methods in HTS: selecting the best from the rest.  
Nat Rev Drug Discov. 2004; 3(2):125-135

## Kinases

- Arner P, Pettersson A, Mitchell PJ, Dunbar JD, Kharitonov A, and Ryden M  
FGF21 attenuates lipolysis in human adipocytes – a possible link to improved insulin sensitivity.  
FEBS Lett. 2008; 582(12):1725-1730
- Binder C, Lafayette A, Archibeque I, Sun Y, Plewa C, Sinclair A, and Emkey R  
Optimization and utilization of the *SureFire*® phospho-STAT5 assay for a cell-based screening campaign.  
Assay and Drug Dev Technol. 2008; 6(1):27-37
- Boldyreff B, Rasmussen TL, Jensen HH, Cloutier A, Beaudet L, Roby P, and Issinger OG  
Expression and purification of PI3 kinase alpha and development of an ATP depletion and an AlphaScreen PI3 kinase activity assay.  
J Biomol Screen. 2008; 13(10):1035-1040

Burns S, Travers J, Collins I, Rowlands MG, Newbatt Y, Thompson N, Garrett MD, Workman P, and Aherne W  
Identification of small-molecule inhibitors of protein kinase B (PKB/AKT) in an AlphaScreen high-throughput screen.  
J Biomol Screen. 2006; 11(7):822-827

Chong MP, Barritt GJ, and Crouch MF  
Insulin potentiates EGFR activation and signaling in fibroblasts.  
Biochem Biophys Res Commun. 2004; 322(2):535-541

Delom F and Chevet E  
Phosphoprotein analysis: from proteins to proteomes.  
Proteome Sci. 2006; 4:15.

Eglen RM and Reisine T  
The current status of drug discovery against the human kinome.  
Assay Drug Dev Technol. 2009; 7(1):22-43. Review

Emery B, Merson TD, Snell C, Young KM, Ernst M, and Kilpatrick TJ  
SOCS3 negatively regulates LIF signaling in neural precursor cells.  
Mol Cell Neurosci. 2006; 31(4):739-747

Guenat S, Rouleau N, Biemann C, Bedard J, Maurer F, Allaman-Pillet N, Nicod P, Bielefeld-Sevigny M, Beckmann JS, Bonny C, Bossé R, and Roduit R  
Homogeneous and nonradioactive high-throughput screening platform for the characterization of kinase inhibitors in cell lysates.  
J Biomol Screen. 2006; 11(8):1015-1026

Hay DL  
Measurement of phosphorylated extracellular signal-regulated kinase 1 and 2 in an undergraduate teaching laboratory with AlphaScreen technology.  
Sci Signal. 2009; 2(62):tr1

Li Y, Cummings RT, Cunningham BR, Chen Y, and Zhou G  
Homogeneous assays for adenosine 5'-monophosphate-activated protein kinase.  
Anal Biochem. 2003; 321(2):151-156

Nishizuka Y  
Intracellular signaling by hydrolysis of phospholipid and activation of protein kinase C.  
Science. 1992; 258(5082):607-614. Review

Roduit R and Schorderet DF  
MAP kinase pathways in UV-induced apoptosis of retinal pigment epithelium ARPE19 cells.  
Apoptosis. 2008; 13(3):343-353

Smotrov N, Mathur A, Kariv I, Moxham CM, and Bays N  
Development of a cell-based assay for measurement of c-Met phosphorylation using AlphaScreen technology and high-content imaging analysis.  
J Biomol Screen. 2009; 14(4):404-411

Stokka AJ, Gesellchen F, Carlson CR, Scott JD, Herberg FW, and Taskén K  
Characterization of A-kinase anchoring disruptors using a solution-based assay.  
Biochem J. 2006; 400(3):493-499

Von Leoprechting A, Kumpf R, Menzel S, Reulle D, Griebel R, Valler MJ, and Buttner FH  
Miniaturization and validation of a high-throughput serine kinase assay using the AlphaScreen platform.  
J Biomol Screen. 2004; 9(8):719-725

Warner G, Illy C, Pedro L, Roby P, and Bossé R  
AlphaScreen kinase HTS platforms.  
Curr Med Chem. 2004; 11(6):721-730

## Lipid:Protein Interactions

Jerga A, Miller DJ, White SW, and Rock CO  
Molecular determinants for interfacial binding and conformational change in a soluble diacylglycerol kinase.  
J Biol Chem. 2009; 284(11):7246-7254

## Nuclear Receptors

Glickman JF, Wu X, Mercuri R, Illy C, Bowen BR, He Y, and Sills M  
A comparison of AlphaScreen, TR-FRET and TRF as assay methods for FXR nuclear receptors.  
J Biomol Screen. 2002; 7(1):3-10

Rouleau N, Turcotte S, Mondou MH, Roby P, and Bossé R  
Development of a versatile platform for nuclear receptor screening using AlphaScreen.  
J Biomol Screen. 2003; 8(2):191-197

Wagstaff K and Jans D  
Intramolecular masking of nuclear localization signals: analysis of importin binding using a novel AlphaScreen-based method.  
Anal Biochem. 2006; 348(1):49-56

Wu X, Glickman JF, Bowen BR, and Sills MA  
Comparison of assay technologies for a nuclear receptor assay screen reveals differences in the sets of identified functional antagonists.  
J Biomol Screen. 2003; 8(4):381-392

Xu HE, Stanley TB, Montana VG, Lambert MH, Shearer BG, Cobb JE, McKee DD, Galardi CM, Plunket KD, Nolte RT, Parks DJ, Moore JT, Kliewer SA, Willson TM, and Stimmel JB  
Structural basis for antagonist-mediated recruitment of nuclear co-repressors by PPAR.  
Nature. 2002; 415: 813-817

## Proteases

Hamilton AC, Inglese J, and Ferrer M  
A PDZ domain-based assay for measuring HIV protease activity: assay design considerations.  
Protein Sci. 2003; 12(3):458-467

Peppard J, Glickman F, He Y, Hu SI, Doughty J, and Goldberg R  
Development of a high-throughput screening assay for inhibitors of aggrecan cleavage using luminescent oxygen channeling (AlphaScreen).  
J Biomol Screen. 2003; 8(2):149-156

## Proteases and Other Enzymatic Assays

Marchand C, Lea WA, Jadhav A, Dexheimer TS, Austin CP, Inglese J, Pommier Y, and Simeonov A  
Identification of phosphotyrosine mimetic inhibitors of human tyrosyl-DNA phosphodiesterase I by a novel AlphaScreen high-throughput assay.  
Mol Cancer Ther. 2009; 8(1):240-248

## Protein:Protein Interactions

Harndahl M, Justesen S, Lamberth K, Røder G, Nielsen M, and Buus S  
Peptide binding to HLA class I molecules: homogenous, high-throughput screening, and affinity assays.  
J Biomol Screen. 2009; 14(2):173-180

Hou Y, McGuinness DE, Prongay AJ, Feld B, Ingravallo P, Ogert RA, Lunn CA, and Howe JA  
Screening for antiviral inhibitors of the HIV integrase-LEDGF/p75 interaction using the AlphaScreen luminescent proximity assay.  
J Biomol Screen. 2008; 13(5):406-414

Jerga A and Rock CO  
Acyl-acyl carrier protein regulates transcription of fatty acid biosynthetic genes via the FabT repressor in *Streptococcus pneumoniae*.  
J Biol Chem. 2009; 284(23):15364-15368

Justesen S, Harndahl M, Lamberth K, Nielsen LL, and Buus S  
Functional recombinant MHC class II molecules and high-throughput peptide-binding assays.  
Immunome Res. 2009; 5:2

Kadkhodayan S, Elliott L, Mausisa G, Wallweber A, Deshayes K, Feng B, and Fairbrother W  
Evaluation of assay technologies for the identification of protein-peptide interaction antagonists.  
Assay Drug Dev Technol. 2007; 5(4):501-513

Kota S, Coito C, Mousseau G, Lavergne JP, and Strosberg AD  
Peptide inhibitors of hepatitis C virus core oligomerization and virus production.  
J Gen Virol. 2009; 90(Pt 6):1319-1328

Mohamed MR, Rahman MM, Lanchbury JS, Shattuck D, Neff C, Dufford M, van Buuren N, Fagan K, Barry M, Smith S, Damon I, and McFadden G  
Proteomic screening of variola virus reveals a unique NF-kappaB inhibitor that is highly conserved among pathogenic orthopoxviruses.  
Proc Natl Acad Sci U S A. 2009; 106(22):9045-9050

Moll D, Prinz A, Gesellchen F, Drewianka S, Zimmermann B, and Herberg FW  
Biomolecular interaction analysis in functional proteomics.  
J Neural Transm. 2006; 113:1015-1032

Rouleau N, Wang J, Karras L, Andrews E, Bielefeld-Sevigny M, and Chen Y  
Highly sensitive assays for SUMOylation and small ubiquitin-like modifier-dependent protein-protein interactions.  
Anal Biochem. 2008; 375(2):364-366

Sehr P, Pawlita M, and Lewis J  
Evaluation of different glutathione S-transferase-tagged protein captures for screening E6/E6AP interaction inhibitors using AlphaScreen.  
J Biomol Screen. 2007; 12(4):560-567

Shan L, Pang L, Zhang R, Murgolo NJ, Lan H, and Hedrick JA  
PCSK9 binds to multiple receptors and can be functionally inhibited by an EGF-A peptide.  
Biochem Biophys Res Commun. 2008; 375(1):69-73

Shih HH, Zhang S, Cao W, Hahn A, Wang J, Paulsen JE, and Harnish DC  
CRP is a novel ligand for the oxidized LDL receptor LOX-1.  
Am J Physiol Heart Circ Physiol. 2009; 296(5):H1643-H1650

Uehara Y, Mochizuki M, Matsuno K, Haino T, and Asai A  
Novel high-throughput screening system for identifying STAT3-SH2 antagonists.  
Biochem Biophys Res Commun. 2009; 380(3):627-631

Yi F and Regan L  
A novel class of small molecule inhibitors of Hsp90.  
ACS Chem Biol. 2008; 3(10):645-654

Yi F, Zhu P, Southall N, Inglese J, Austin CP, Zheng W, and Regan L  
An AlphaScreen-based high-throughput screen to identify inhibitors of Hsp90-cochaperone interaction.  
J Biomol Screen. 2009; 14(3):273-281

## Single Nucleotide Polymorphism (SNP) Detection

Beaudet L, Bedard J, Breton B, Mercuri RJ, and Budarf ML  
Homogeneous assays for single-nucleotide polymorphism typing using AlphaScreen.  
Genome Res. 2001; 11(4):600-608

Tsuchihashi Z and Dracopoli NC  
Progress in high-throughput SNP genotyping methods.  
Pharmacogenomics J. 2002; 2(2):103-110. Review

## Ubiquitin Ligases

Kus B, Gajadhar A, Stanger K, Cho R, Sun W, Rouleau N, Lee T, Chan D, Wolting C, Edwards A, Bossé R, and Rotin D  
A high-throughput screen to identify substrates for the ubiquitin ligase Rsp5.  
J Biol Chem. 2005; 280(33):29470-29478